



Supplementary material to Electrochemical determination of paracetamol and chlorpheniramine using gold nanoparticles/reduced graphene oxide modified glassy carbon electrode

Huynh Van Chung^{1,2}, Nguyen Hai Phong¹, Dao Thi Cam Minh³, Ho Xuan Anh Vu¹,
Pham Khac Lieu¹, Nguyen Thi Thanh Huyen⁴, Trinh Ngoc Dat⁵, Doan Manh Dung⁶, Do
Thi Duyen⁷, Dinh Quang Khieu¹

¹University of Sciences, Hue University, Hue City, Vietnam

²Faculty of Pharmacy, Buon Ma Thuot Medical University, Buon Ma Thuot City, Vietnam

³University of Medicine and Pharmacy, Hue University, Hue City, Vietnam

⁴Institute of Research and Development, Duy Tan University, Danang City, Vietnam

⁵University of Education and Science, The University of Danang, Danang City, Vietnam

⁶Institute of Biotechnology and Environment, Tay Nguyen University, Buon Ma Thuot, Vietnam

⁷Institute of Materials Science, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet,
Cau Giay, Ha Noi city, Vietnam

J. Electrochem. Sci. Eng. 00(0) (2025) 2841; <http://dx.doi.org/10.5599/jese.2841>




Figure S1. a) SqW-ASV curves of AuNPs-rGO-GCE and b) I_p variation (standard deviations marked) of PAR and CPM at different accumulation potentials ($t_{acc} = 60$ s, $f = 20$ Hz, $\Delta E = 25$ mV, $U_{step} = 6$ mV, $v = 25$ mV s⁻¹, $C_{PAR} = 28$ μ M, $C_{CPM} = 54$ μ M, BR buffer 0.05 M, pH 7)




Figure S2. a) SQW-ASV curves of AuNPs-rGO-GC and b) I_p variation (standard deviations marked) of PAR and CPM with different accumulation times ($E_{acc} = -900$ mV, $f = 20$ Hz, $\Delta E = 25$ mV, $U_{Step} = 6$ mV, $v = 25$ mV s⁻¹, $C_{PAR} = 28$ μM , $C_{CPM} = 54$ μM , BR buffer 0.05 M, pH 7)




Figure S3. a) SQW-ASV curves of AuNPs-rGO-GCE and b) I_p variation (standard deviations marked) of PAR and CPM with wave frequencies from 5 to 50 Hz corresponding to sweep rate from 30 to 300 mV s⁻¹ ($E_{acc} = -900$ mV, $t_{acc} = 30$ s, $\Delta E = 25$ mV, $U_{Step} = 6$ mV, $v = 25$ mV s⁻¹, $C_{PAR} = 28$ μM , $C_{CPM} = 54$ μM , BR buffer 0.05 M, pH 7)




Figure S4. a) SqW-ASV curves of AuNPs-rGO-GCE and b) I_p and c) E_p variations (standard deviations marked) of PAR and CPM with different pulse amplitudes. ($E_{acc} = -900$ mV, $t_{acc} = 30$ s, $f = 20$ Hz, $U_{Step} = 6$ mV, $v = 120$ mV s⁻¹, $C_{PAR} = 28$ μM , $C_{CPM} = 54$ μM , BR buffer 0.05 M, pH 7)




Figure S5. SQW-ASV curves of AuNPs-rGO in 0.05 M BR buffer, pH 7, containing PAR, CPM, and uric acid (UA), caffeine (CAF) and ascorbic acid (AA). ($E_{acc} = -900$ mV, $t_{acc} = 30$ s, $\Delta E = 30$ mV, $U_{Step} = 6$ mV, $v = 120$ mV s⁻¹, $C_{PAR} = C_{CPM} = 28$ μM , $C_{UA} = C_{AA} = C_{CAF} = 112$ μM)




Figure S6. SQW-ASV curves of of AuNPs-rGO-GCE in which solution containing PAR, CPM, and various inorganic compounds, each at a concentration of 1120 μM . ($E_{\text{acc}} = -900 \text{ mV}$, $t_{\text{acc}} = 30 \text{ s}$, $\Delta E = 30 \text{ mV}$, $U_{\text{Step}} = 6 \text{ mV}$, $v = 120 \text{ mV s}^{-1}$, $C_{\text{PAR}} = C_{\text{CPM}} = 28 \mu\text{M}$ BR buffer 0.05 M, pH 7).